

**VIRUCIDAL EFFICACY<sup>+</sup>***Inactivation of enveloped viruses as SARS-CoV-2 in domestic laundry processes***SARS-CoV-2**

The novel human coronavirus, SARS-CoV-2 has become a global health concern in 2020, causing COVID-19, a severe respiratory tract infection in humans. Only few data are so far available for SARS-CoV-2, but representative data exists for related coronaviruses causing SARS or MERS. The novel coronavirus, SARS-CoV-2, is transmitted via droplet (coughing, sneezing) or hand- and surface contact.

**Enveloped versus non-enveloped**

An important aspect for the inactivation of viruses is the discrimination between enveloped and non-enveloped viruses. Non-enveloped viruses as e.g. Norovirus are in general more difficult to inactivate and require higher disinfectant concentrations and temperatures, whereas the enveloped SARS-CoV-2 is easier to inactivate in a washing process (Sidwell & Dixon, 1969; Gerba, 2001; Gerba & Kennedy, 2007; Gerhardts et al., 2009; Heinzel et al., 2010).

**Textile hygiene**

Textile fabrics may play a role in the spread of infectious viral diseases (Bloomfield, et al., 2008; Gerhardts et al., 2012). It is therefore crucial that a washing process reduces the virus load on textiles to a safe level.

**Washing process**

Several studies showed that enveloped viruses are fully inactivated at 20 to 30 °C with an active oxygen bleach containing detergent (powder detergent for white laundry) (Heinzel et al., 2010; Gerhardts et al., 2016). However, without detergent some active virus particles could be recovered after a 30 °C washing cycle and a carry-over to other laundry items could be observed (Gerhardts et al., 2016).

**Recommendation for SARS-CoV-2**

Even though for enveloped viruses as SARS-CoV-2 a washing temperature of 20 to 30 °C with a bleach-containing detergent seems to be sufficient, it is recommended to use  $\geq 40$  °C with active oxygen bleach containing powder detergent for household washing processes.

**Standards**

No standard is available to assess the virucidal action of either I&I or domestic laundry processes. The closest standards are those for testing disinfectants as the EN 13610 (food and industrial areas), EN 14476 (medical area) or EN 14675 (veterinary area). All standards require a virus reduction of  $\geq 4 \log_{10}$ .

**Testing of laundry processes**

MS2-phage is a model virus for the non-enveloped Norovirus or Hepatitis A virus with comparable resistance against temperature, ozone, UV, peracetic acid and sodium hypochlorite (Allwood, Malik, Hedberg & Goyal, 2003; De Roda Husman et al., 2004; Park, Boston, Kase, Sampson & Sobsey, 2007; Shin & Sobsey, 2003; Morin et al., 2015). Since SARS-CoV-2 is enveloped a lower persistence is expected as for MS2-phage in a washing process. A full inactivation of MS2-phage would indicate also an inactivation of SARS-CoV-2.

## HyWa-Check Biomonitor

Alternatively, also HyWa-Check biomonitor No. 503 (*Staphylococcus arlettae*) or No. 504 (*Enterococcus faecium*) can be used to test the efficiency of a washing process. Enveloped viruses are expected to have an equal to lower resistance towards temperature, surfactants and bleach as bacteria (Bloomfield et al., 2013). The HyWa-Check biomonitor No. 503 and 504 contain gram-positive bacteria with a relatively high tolerance towards temperature and bleach. An inactivation of *Enterococcus faecium* or *Staphylococcus arlettae* is therefore an indicator for a washing process sufficiently effective against enveloped viruses as SARS-CoV-2.

## Swissatest

Please contact our hygiene department for further informations on virucidal efficacy testing of I&I or household laundry processes. Swissatest provides either the possibility to test the virucidal action with MS2 phage or with HyWa-Check biomonitor. Please be aware that delivery time of HyWa-Check biomonitor is 3-4 weeks.

## Contact

Swissatest Testmaterialien AG  
Mövenstrasse 12  
CH-9015 St.Gallen  
Switzerland  
Mail: [info@swissatest.ch](mailto:info@swissatest.ch)  
Phone: +41 71 311 80 55

## References

- Allwood, P. B., Malik, Y. S., Hedberg, C. W., Goyal, S. M. (2003). *Appl Environ Microbiol*, 69, 5707-5710.
- Bloomfield, S., Exner, M., et al. (2008). *Eurosurveillance*, 13, 1-4.
- Bloomfield, S. F., Exner, M., Signorelli, C. & Scott, E. A. (2013). *Int Sci Forum Home Hyg* (October), 1-62.
- De Roda Husman, A. M., Bijkerk, P., Lodder, W. et al. (2004). *Appl Environ Microbiol*, 70, 5089-5093.
- Gerba, C. P. (2001). *J Infect*, 43(1), 92-98.
- Gerba, C. P. & Kennedy, D (2007). *App Env Microbiol*, 73(14), 4425-4428. "
- Gerhardts, A., Wilderer, C., Mucha, H. & Höfer, D. (2009). *Hyg und Medizin*, 34(7/8), 272-281.
- Gerhardts, A., Mucha, H., et al. (2012). *Hygiene & Medizin*, 37, 400-403.
- Gerhardts, A., et al. (2016). *Journal of Biosciences and Medicines*, 4, 111-125.
- Heinzel, M., Kyas, A., Wiede, M., Vreves, R. & Bockmuehl, D.P. (2010). *Int J Hyg Environ Health*, 213(5), 334-337.
- Morin, T. et al (2015). *Journal of Applied Microbiology*, 119, 6555-665.
- Shin, G. A. & Sobsey, M. D. (2003). *Appl Environ Microbiol*, 69, 3975-3978.
- Sidwell, R. & Dixon, G. (1969). *Am Oil Chem Soc*, 46(10), 532-536.
- Park, G. W., Boston, D. M., Kase, J. A., Sampson, M. N., & Sobsey, M. D. (2007). *Appl Environ Microbiol*, 73, 4463-4468.